

## MORPHOLOGICAL FEATURES OF THE SMALL INTESTINAL INJURIES IN THE EARLY PERIOD OF MILD SPINAL BRAIN INJURY

Sidikova Gulrukh Shavkatovna

Department of "Propaedeutics of children's diseases and children's neurology" Bukhara State Medical  
Institute

**Abstract:** *This thesis presents the results of scientific studies that allowed us to assess and study the morphological features of the small intestine of 3-month-old white crossbred rats in the early stages of mild spinal cord injury.*

**Keywords:** *proximal and distal parts of the small intestine, villi, atrophy, hemorrhagic hemorrhage.*

**Abstract:** *This thesis presents the results of scientific studies that allowed us to evaluate and study the morphological characteristics of the small intestine of 3-month-old white outbred rats in the early stages of mild spinal cord injury.*

**Keywords:** *proximal and distal small intestine, villi, atrophy, hemorrhagic hemorrhage.*

The incidence and prevalence of spinal cord and spinal cord injuries worldwide have increased by almost thirtyfold in the past few years [1,2]. According to WHO, up to now, about 500,000 people suffer spinal cord and spinal cord injuries every year due to various car accidents and accidental events, which ultimately lead to group I and II disability of patients. Because the prognosis for the recovery processes related to sensory and motor functions after spinal cord injury is unsatisfactory [3].

The extent of dysfunctional changes in the digestive system and other systems in patients with spinal cord and spinal cord injuries largely depends on the location and severity of the injury, among which complications in the small intestine are of particular importance. Timely and successful treatment and prevention of complications in the small intestine due to spinal cord and spinal cord injuries allows preserving the functions of these organs and improving the quality of life of patients, as well as saving the lives of victims by reducing digestive complications. Data on morphological changes in digestive organs after spinal cord injury, especially changes in the small intestine, have been poorly studied, which shows that complications in this organ are not only a medical, but also an economic and social problem, and finding a solution to them is an urgent task.

**Materials and methods:** The experiments were conducted on 12 3-month-old white male and female rats born in vivarium conditions. In order to determine the morphological parameters of the structure of the small intestine in the early period of spinal cord injury, the following 2 large groups were formed:

1. Control group (n=5)
2. Research group, i.e., mild spinal cord injury but not treated with drugs (n=7)

In the study, mild spinal cord injury was inflicted on white male rats using a specially developed model using the "fall from a height" method. Then, the animals of the experimental group were killed by decapitation under light isoflurane anesthesia and the small intestine was isolated. Histological preparations were prepared from the excised

organ and stained with hematoxylin and eosin. Then, corresponding histological images were taken from the preparations.

Research results: The studies showed that the body weight of three-month-old white rats in the 2nd (study) group ranged from 120 g to 158 g, with an average of 144 g. The early period of spinal cord injury, usually 14 days after injury, is characterized by the presence of inflammation and secondary degenerative changes in the wall of the small intestine due to disruption of innervation by the nervous system.

When the small intestine of rats in the 2nd study group was stained with hematoxylin and eosin, we can see a decrease in the height of the villi and the area they occupy, and a flattening of their surface. This condition sharply reduces the surface area of absorption of nutrients from the small intestine. We can witness the appearance of signs of hypertrophy of the intestinal muscle layer. If this condition continues for a long time, it leads to chronic disorders of intestinal motility.

Conclusion: Thus, in the 2nd study group, 3-month-old white purebred rats that received mild spinal cord injury but were not treated with drugs, histologically, various pathological changes were noted in the wall of the proximal and distal parts of the small intestine.

#### REFERENCES:

1. Devivo M.J. Epidemiology of traumatic spinal cord injury: trends and future implications // *Spinal Cord*. 2012 May;50(5):365-72.
2. Nemunaitis, G. Early Predictors of Functional Outcome After Trauma / G. Nemunaitis, M.J. Roach, J. Claridge et al. // *J injury function rehabil.* - 2015. - Vol. 24. - S 1482-1934.
3. Sminkey L. Международный обзор травматизма позвоночника // ВООЗ, Женева – Презентация